

英文原著

Ikeda H, Shiku H. Antigen-receptor gene-modified T cells for treatment of glioma. *Glioma: Immunotherapeutic approaches*. Chapter 17. 2011.

Tobinai K, Yamaguchi M. Treatment of extranodal NK/T cell lymphoma, nasal type. UpToDate™ (Last major updated date: Dec 23, 2011)

Miyazaki K, Yamaguchi M, Suzuki R, Kobayashi Y, Maeshima AM, Niitsu N, Ennishi D, Tamaru JI, Ishizawa K, Kashimura M, Kagami Y, Sunami K, Yamane H, Nishikori M, Kosugi H, Yujiri T, Hyo R, Katayama N, Kinoshita T, Nakamura S. CD5-positive diffuse large B-cell lymphoma: a retrospective study in 337 patients treated by chemotherapy with or without rituximab. *Ann Oncol*. 22: 1601-1607, 2011.

Umino A, Nakagawa M, Utsunomiya A, Tsukasaki K, Taira N, Katayama N, Seto M. Clonal evolution of adult T cell leukemia/lymphoma takes place in the lymph nodes. *Blood*. 117(20):5473-5478, 2011.

Jankowska AM, Makishima H, Tiu RV, Szpurka H, Huang Y, Traina F, Visconte V, Sugimoto Y, Prince C, O'Keefe C, Hsi ED, List A, Sekeres MA, Rao A, McDevitt MA, Maciejewski JP. Mutational spectrum analysis of chronic myelomonocytic leukemia includes genes associated with epigenetic regulation: UTX, EZH2, and DNMT3A. *Blood*. 118(14):3932-3941, 2011

Suzuki R, Yamaguchi M, Izutsu K, Yamamoto G, Takada K, Harabuchi Y, Isobe Y, Gomyo H, Koike T, Okamoto M, Hyo R, Suzumiya J, Nakamura S, Kawa K, Oshimi K. Prospective measurement of Epstein-Barr virus-DNA in plasma and peripheral blood mononuclear cells of extranodal NK/T-cell lymphoma, nasal type. *Blood* 118:6018-6022, 2011

Lesokhin AM, Hohl TM, Kitano S, Cortez C, Hirschhorn-Cymerman D, Avogadri F, Rizzuto GA, Lazarus JJ, Eric G, Pamer EG, Houghton AN, Merghoub T, Wolchok JD. **Monocytic** CCR2⁺ Myeloid Derived Suppressor Cells Promote Immune Escape By Limiting

Activated CD8 T Cell Infiltration Into The Tumor Microenvironment. *Cancer Res* 72(4):876-886, 2011

Shirakura Y, Mizuno Y, Wang L, Imai N, Amaike C, Sato E, Ito M, Nukaya I, Mineno J, Takesako K, Ikeda H, Shiku H. T-cell receptor gene therapy targeting melanoma-associated antigen-A4 inhibits human tumor growth in non-obese diabetic/SCID/gc^{null} mice. *Cancer Sci*. Epub ahead of print, 2011

Tsuchikawa T, Ikeda H, Cho Y, Miyamoto M, Shichinohe T, Hirano S, Kondo S. Association of CD8⁺ T cell infiltration in esophageal carcinoma lesions with HLA class I antigen expression and survival. *Clin. Exp. Immunol.* 164(1):50-56, 2011

Takahashi N, Wakita H, Miura M, Scott SA, Nishii K, Masuko M, Sakai M, Maeda Y, Ishige K, Kashimura M, Fujikawa K, Fukazawa M, Katayama T, Monma F, Narita M, Urase F, Furukawa T, Miyazaki Y, Katayama N, Sawada K. Correlation between imatinib pharmacokinetics and clinical response in Japanese patients with chronic phase chronic myeloid leukemia. *Clin Pharmacol Ther.* 88(6):809-813, 2010.

Iwamoto T, Monma F, Fujieda A, Nakatani K, Katayama N, Okuda M. Hepatic drug interaction between tacrolimus and lansoprazole in a bone marrow transplant patient receiving variconazole and harboring CYP2C19 and CYP3A5 heterozygous mutations. *Clin Ther.* 33(8):1077-1080, 2011.

Tano S, Tanaka K, Iguchi T, Nishikawa K, Takayama R, Ninomiya K, Inoue H, Katsurahara M, Horiki N, Takei Y. Large retention cyst with chondroid metaplasia in esophagus. *Endoscopy* 43 Suppl 2 UCTN:E262-263, 2011.

Nagai H, Ogura M, Kusumoto S, Takahashi N, Yamaguchi M, Takayama N, Kinoshita T, Motoji T, Ohyashiki K, Kosugi H, Matsuda S, Ohnishi K, Omachi K, Hotta T. Cladribine combined with rituximab (R-2-CdA) therapy is an effective salvage therapy in relapsed or refractory indolent B-cell non-Hodgkin lymphoma. *Eur J Haematol* 86:117-123, 2011

Masuya M, Nakamura S, Yukimoto H, Miyata E, Ino K, Liu B, Suzuki K, Ohishi K, Katayama N. Ly6C(+) monocytes are extrahepatic precursors of hepatic stellate cells in the injured liver of mice. *Exp Hematol.* 39(9):934-946, 2011

Iwamura K, Kato T, Miyahara Y, Naota H, Mineno J, Ikeda H, Shiku H. siRNA-mediated silencing of PD-1 ligands enhances tumor-specific human T cell effector functions. *Gene Ther.* Epub ahead of print, 2011

Afable MG 2nd, Shaik M, Sugimoto Y, Elson P, Clemente M, Makishima H, Sekeres MA, Lichtin A, Advani A, Kalaycio M, Tiu RV, O'Keefe CL, Maciejewski JP. Efficacy of rabbit anti-thymocyte globulin in severe aplastic anemia. *Haematologica.* 96(9):1269-1275, 2011

Okano H, Isono Y, Tanaka H, Ishihara Y, Matsusaki S, Aoki M, Sase T, Saitou T, Mukai K, Nishimura A, Takemoto M, Yokote K. Primary liver tumor associated with Werner syndrome (adult progeria). *Hepatology Research* 41:1260-5, 2011.

Ito-Habe N, Wada H, Matsumoto T, Ohishi K, Toyoda H, Ishikawa E, Nomura S, Komada Y, Ito M, Nobori T, Katayama N. Elevated Von Willebrand factor propeptide for the diagnosis of thrombotic microangiopathy and for predicting a poor outcome. *Int J Hematol.* 93(1):47-52, 2011

Okano H, Saitou T, Sase T. Bleeding from a Gastric Hyperplastic Polyp during Anticoagulation Therapy. *Internal Medicine* 50:1349, 2011

Kaneko T, Wada H. Diagnostic criteria and laboratory tests for disseminated intravascular coagulation. *J Clin Exp Hematop.* 51(2):67-76, 2011.

Watanabe T, Tobinai K, Shibata T, Tsukasaki K, Morishima Y, Maseki N, Kinoshita T, Suzuki T, Yamaguchi M, Ando K, Ogura M, Taniwaki M, Uike N, Takeuchi K, Nawano S, Terauchi T, Hotta T. Phase II/III study of R-CHOP-21 versus R-CHOP-14 for untreated indolent B-cell non-Hodgkin's lymphoma: JCOG 0203 trial. *J Clin Oncol* 29: 3990-3998, 2011

Yamaguchi M, Kwong Y-L, Kim WS, Maeda Y, Hashimoto C, Suh C, Izutsu K, Ishida F, Isobe Y, Sueoka E, Suzumiya J, Kodama T, Kimura H, Hyo R, Nakamura S, Oshimi K, Suzuki R. Phase II study of SMILE chemotherapy for newly diagnosed stage IV, relapsed, or refractory extranodal natural killer (NK)/T-cell lymphoma, nasal type: the NK-Cell Tumor Study Group study. *J Clin Oncol* 29: 4410-4416, 2011

Omoto Y, Yamanaka K, Tokime K, Kitano S, Kakeda M, Akeda T, Kurokawa I, Gabazza EC, Tsutsui H, Katayama N, Yamanishi K, Nakanishi K, Mizutani H. Granzyme B is a novel interleukin-18 converting enzyme. *J Dermatol Sci.* 59(2):129-135, 2010.

Horiki N, Omata F, Uemura M, Suzuki S, Ishii N, Fukuda K, Fujita Y, Ninomiya K, Tano S, Katurahara M, Tanaka K, Gabazza EC, Takei Y. Risk for local recurrence of early gastric cancer treated with piecemeal endoscopic mucosal resection during a 10-year follow-up period. *Surg Endosc* 26(1):72-78, 2012.

Takemitsu T, Wada H, Hatada T, Ohmori Y, Ishikura K, Takeda T, Sugiyama T, Yamada N, Maruyama K, Katayama N, Isaji S, Shimpo H, Kusunoki M, Nobori T. Prospective evaluation of three different diagnostic criteria for disseminated intravascular coagulation. *Thromb Haemost.* 105(1):40-44, 2011.

Nakamura A, Abe K, Masuya M, Imai S, Ohishi K, Mori Y, Kojima T, Wada H, Katayama N, Nobori T. Efficiency of diversion of the first aliquot of blood and prestorage leukoreduction for preventing bacterial contamination in red blood cell concentrates assessed using a rapid polymerase chain reaction-based bacterial detection system. *Transfus Med.* 21(6):365-370, 2011