

英文原著

Shirakura Y, Mizuno Y, Wang L, Imai N, Amaike C, Sato E, Ito M, Nukaya I, Mineno J, Takesako K, Ikeda H, Shiku H.

T-cell receptor gene therapy targeting melanoma-associated antigen-A4 inhibits human tumor growth in non-obese diabetic/SCID/ γ c null mice.

Cancer Sci. 103(1):17-25, 2012

Iwamura K, Kato T, Miyahara Y, Naota H, Mineno J, Ikeda H, Shiku H.

si-RNA-mediated silencing of PD-1 ligands enhances tumor-specific human T-cell effector functions.

Gene Ther. 19(10):959-966, 2012

Noguchi T, Kato T, Wang L, Maeda Y, Ikeda H, Sato E, Knuth A, Gnjatic S, Ritter G, Sakaguchi S, Old LJ, Shiku H, Nishikawa H.

Intracellular tumor-associated antigens represent effective targets for passive immunotherapy.

Cancer Res. 72(7):1672-1682, 2012

Ikeda H.

Gene-modified lymphocytes: from caution to promise for effective cancer immunotherapy.

Immunotherapy. 4(3):241-244, 2012

Ikeda H, Shiku H.

Antigen-receptor gene-modified T cells for treatment of glioma.

Adv Exp Med Biol. 746:202-215, 2012

Okamoto S, Amaishi Y, Goto Y, Ikeda H, Fujiwara H, Kuzushima K, Yasukawa M, Shiku H, Mineno J.

A Promising Vector for TCR Gene Therapy: Differential Effect of siRNA, 2A Peptide, and Disulfide Bond on the Introduced TCR Expression.

Mol Ther Nucleic Acids. 1:e63, 2012

Nakamori Y, Liu B, Ohishi K, Suzuki K, Ino K, Matsumoto T, Masuya M, Nishikawa H, Shiku H, Hamada H, Katayama N.

Human bone marrow stromal cells simultaneously support B and T/NK lineage development from human haematopoietic progenitors: a principal role for flt3 ligand in lymphopoiesis.

Br J Haematol. 157(6):674-686, 2012

Suzuki K, Ono R, Ohishi K, Masuya M, Kataoka I, Liu B, Nakamori Y, Ino K, Monma F, Hamada H, Kitamura T, Katayama N, Nosaka T.

IKAROS isoform 6 enhances BCR-ABL1-mediated proliferation of human CD34+ hematopoietic cells on stromal cells.

Int J Oncol. 40(1):53-62, 2012.

Habe K, Wada H, Ito-Habe N, Hatada T, Matsumoto T, Ohishi K, Maruyama K, Imai H, Mizutani H, Nobori T.

Plasma ADAMTS13, von Willebrand Factor (VWF) and VWF Propeptide Profiles in Patients with DIC and Related Diseases.

Thromb Res. 129(5):598-602, 2012.

Sakata K, Matsumoto T, Wada H, Kumamoto T, Iwamoto S, Ohishi K, Takahashi J, Katayama N, Komada Y.

A case of three-year-old boy with severe haemophilia A on prophylaxis identified with a chronic subdural haematoma 2 years after the first episode of acute subdural bleeding. Haemophilia 18(2), e1-2, 2012

Hamada Y, Tanaka K, Tano S, Katsurahara M, Kosaka R, Noda T, Aoki M, Toyoda H, Takei Y, Katayama N.

Usefulness of endoscopic submucosal dissection for the treatment of rectal carcinoid tumors.

Eur J Gastroenterol Hepatol. 24(7):770-774, 2012

Kosaka R, Tanaka K, Tano S, Takayama R, Nishikawa K, Hamada Y, Toyoda H, Ninomiya K, Katsurahara M, Inoue H, Horiki N, Katayama N, Takei Y.

Magnifying endoscopy for diagnosis of residual/local recurrent gastric neoplasms after previous endoscopic treatment.

Surg Endosc. 26(8):2299-2305, 2012.

Tawara I, Sun Y, Liu C, Toubai T, Nieves E, Evers R, Alrubaie M, Mathewson N, Tamaki H, Reddy P.

Donor- but not host-derived interleukin-10 contributes to the regulation of experimental graft-versus-host disease.

J Leukoc Biol. 91(4):667-675, 2012

Toubai T, Tawara I, Sun Y, Liu C, Nieves E, Evers R, Friedman T, Korngold R, Reddy P.

Induction of acute GVHD by sex-mismatched H-Y antigens in the absence of functional radiosensitive host hematopoietic-derived antigen presenting cells.

Blood 119(16):3844-3853, 2012

Tawara I, Sun Y, Lewis EC, Toubai T, Evers R, Nieves E, Azam T, Dinarello CA, Reddy P.

Alpha-1-antitrypsin monotherapy reduces graft-versus-host disease after experimental allogeneic bone marrow transplantation.

Proc Natl Acad Sci U.S.A. 109(2):564-569, 2012

Tada K, Kim SW, Asakura Y, Hiramoto N, Yakushijin K, Kurosawa S, Tajima K, Mori S, Heike Y, Tanosaki R, Maeshima AM, Taniguchi H, Furuta K, Kagami Y, Matsuno Y, Tobinai K, Takaue Y, Fukuda T.

Comparison of outcomes after allogeneic hematopoietic stem cell transplantation in patients with follicular lymphoma, diffuse large B-cell lymphoma associated with follicular lymphoma, or de novo diffuse large B-cell lymphoma.

Am J Hematol. 87(8):770-775, 2012

Yamaguchi M.

Current and future management of NK/T-cell lymphoma based on clinical trials.

Int J Hematol. 96(5):562-571, 2012

Yamaguchi M, Tobinai K, Oguchi M, Ishizuka N, Kobayashi Y, Isobe Y, Ishizawa K, Maseki N, Itoh K, Usui N, Wasada I, Kinoshita T, Hotta T, Tsukasaki K, Oshimi K. Concurrent chemoradiotherapy for localized nasal natural killer/T-cell lymphoma: an updated analysis of the Japan Clinical Oncology Group Study JCOG0211. *J Clin Oncol.* 30(32):4044-4046, 2012

Ito Y, Kimura H, Maeda Y, Hashimoto C, Ishida F, Izutsu K, Fukushima N, Isobe Y, Takizawa J, Hasegawa Y, Kobayashi H, Okamura S, Kobayashi H, Yamaguchi M, Suzumiya J, Hyo R, Nakamura S, Kawa K, Oshimi K, Suzuki R. Pretreatment EBV-DNA copy number is predictive of response and toxicities to SMILE chemotherapy for extranodal NK/T-cell lymphoma, nasal type. *Clin Cancer Res.* 18(15):4183-4190, 2012

Jerez A, Sugimoto Y, Makishima H, Verma A, Jankowska AM, Przychodzen B, Visconte V, Tiu RV, O'Keefe CL, Mohamedali AM, Kulasekararaj AG, Pellagatti A, McGraw K, Muramatsu H, Moliterno AR, Sekeres MA, McDevitt MA, Kojima S, List A, Boulwood J, Mufti GJ, Maciejewski JP. Loss of heterozygosity in 7q myeloid disorders: clinical associations and genomic pathogenesis. *Blood* 119(25):6109-6117, 2012

Sugimoto Y, Sekeres MA, Makishima H, Traina F, Visconte V, Jankowska A, Jerez A, Szpurka H, O'Keefe CL, Guinta K, Afable M, Tiu R, McGraw KL, List AF, Maciejewski J. Cytogenetic and molecular predictors of response in patients with myeloid malignancies without del[5q] treated with lenalidomide. *J Hematol Oncol.* 5(1):4, 2012

Makishima H, Sugimoto Y, Szpurka H, Clemente MJ, Ng KP, Muramatsu H, O'Keefe C, Sauntharajah Y, Maciejewski JP. CBL mutation-related patterns of phosphorylation and sensitivity to tyrosine kinase inhibitors.

Leukemia 26(7):1547-1554, 2012

Hama A, Muramatsu H, Makishima H, Sugimoto Y, Szpurka H, Jasek M, O'Keefe C, Takahashi Y, Sakaguchi H, Doisaki S, Shimada A, Watanabe N, Kato K, Kiyoi H, Naoe T, Kojima S, Maciejewski JP.

Molecular lesions in childhood and adult acute megakaryoblastic leukaemia.

Br J Haematol. 156(3):316-325, 2012

Hirschhorn-Cymerman D, Budhu S, Kitano S, Liu C, Zhao F, Zhong H, Lesokhin AM, Avogadri-Connors F, Yuan J, Houghton AN, Merghoub T, Wolchok JD.

Induction of potent tumoricidal function in CD4+ T cells is associated with concomitant memory and terminally differentiated phenotype in mice and humans.

J Exp Med. 209(11):2113-2126, 2012

Postow MA, Callahan MK, Barker CA, Yamada Y, Yuan J, Kitano S, Mu Z, Rasalan T, Adamow M, Ritter E, Sedrak C, Jungbluth AA, Chua R, Yang AS, Roman RA, Rosner S, Benson B, Allison JP, Lesokhin AM, Gnjatic S, Wolchok JD.

Immunologic correlates of the abscopal effect in a patient with melanoma.

N Engl J Med. 366(10):925-931, 2012

Lesokhin AM, Hohl TM, Kitano S, Cortez C, Hirschhorn-Cymerman D, Avogadri F, Rizzuto GA, Lazarus JJ, Pamer EG, Houghton AN, Merghoub T, Wolchok JD.

Monocytic CCR2+ myeloid-derived suppressor cells promote immune escape by limiting activated CD8 T-cell infiltration into the tumor microenvironment.

Cancer Res. 72(4):876-886, 2012

Suzuki K, Nakamura A, Fujieda A, Nakase K, Katayama N.

Pulmonary infection caused by *Exophiala dermatitidis* in a patient with multiple myeloma: A case report and a review of the literature.

Medical Mycology Case Reports 1(1): 95-98, 2012

Kawakami K, Ito R, Tono Y, Yazaki A, Murata T, Baba Y, Uchiyama T, Ito M.

Orbital inflammatory lesion as an initial manifestation of systemic nasal type NK/T cell

lymphoma.

J Clin Exp Hematopathol. 52(2):137-139, 2012

Fujiwara A, Kobayashi H, Masuya M, Maruyama M, Nakamura S, Ibata H, Fujimoto H, Ohnishi M, Urawa M, Naito M, Takagi T, Kobayashi T, Gabazza EC, Takei Y, Taguchi O.

Correlation between circulating fibrocytes, and activity and progression of interstitial lung diseases.

Respirology 17(4):693-698, 2012

Nakase K, Suzuki K, Kyo T, Kohara T, Sugawara Y, Katayama N.

Is elevation of the serum β -D-glucan level a paradoxical sign for Trichosporon fungemia in patients with hematologic disorders?

International Journal of Infectious Diseases 16:e2-e4, 2012

Nakase K, Kita K, Kyo T, Tsuji K, Katayama N.

High serum levels of soluble interleukin-2 receptor in acute myeloid leukemia: correlation with poor prognosis and CD4 expression on blast cells.

Cancer Epidemiol. 36(5):e306-e309, 2012

Tobinai K, Yamaguchi M.

Treatment of extranodal NK/T cell lymphoma, nasal type.

UpToDate™